

*C. sagitta*, Gask., is also an *Amycla*. In Mr. Pease's series of specimens, the arrow-pattern gradually passes away.

*Sistrum seminodosum*, Pse., = *elatum*, Blainv., var.

*Mitra brunnea*, Pse., = *Strigatella fuscescens*, Pse., var. (teste H. Cuming).

*M. tessellata*, Pse., = *M. puella*, Rve. (teste H. Cuming).

*Triton crenulatus*, Pse., = *Epidromus antiquatus*, Hds.

3. DESCRIPTIONS OF FOUR NEW SCINTILLÆ AND ONE NEW PYTHINA FROM THE COLLECTION OF MR. H. CUMING. BY G. B. SOWERBY, F.L.S.

(Pl. XXXII.)

SCINTILLA SEMICLAUSA. Pl. XXXII. f. 1, 2. *Testa oblongo-subquadrata, subventricosa, paulo obliquata, succinea, ad marginem ventralem clausa; lateribus vix hiantibus; apice acuminato; latere antico brevior et angustior.*

*Hab.* Borneo (*Mus. H. Cuming*).

Less obtusely angled on the anterior side than *S. succinea*, and deeper than *S. crocea*.

SCINTILLÆ LACTEA. Pl. XXXII. f. 4. *Testa oblique ovalis, alba, clausa, subventricosa, latere postico subextenso.*

*Hab.* Borneo (*Mus. H. Cuming*).

More oblique and less obtuse at the sides than *S. layardi*.

SCINTILLA ROSEA. Pl. XXXII. f. 5, 6. *Testa ovali-oblonga, compressa, rosea, antice subacuta, breviuscula, postice lata rotundata; margine ventrali expanso.*

*Hab.* Lizard Islands (*Mus. H. Cuming*).

More expanded at the ventral margin, and more pointed anteriorly, than *S. pudica*.

SCINTILLA OBLONGA. Pl. XXXII. f. 3. *Testa pallide straminea, oblonga, lateribus subquadratis, marginibus dorsalibus et ventralibus rectis.*

*Hab.* Borneo (*Mus. H. Cuming*).

More quadrately oblong than *S. borneensis* and similar species.

PYTHINA STRIATISSIMA. Pl. XXXII. f. 7. *Testa pallide straminea, minutissime divaricatim striata; complanata, subtrigona, marginibus latero-dorsalibus declivis, margine ventrali rectiusculo, medio vix sinuato.*

*Hab.* Borneo (*Mus. H. Cuming*).